



RESEARCH PROGRESS REPORT SUMMARY

Grant 02686-A: Pattern of Thyroid Function Tests during Recovery from Acute Nonthyroidal Illness

Principal Investigator: Timothy Bolton, DVM

Research Institution: Virginia-Maryland Regional College of Veterinary Medicine

Grant Amount: \$13,792

Start Date: 1/1/2020 **End Date:** 6/30/2022

Progress Report: End-Year 1

Report Due: 12/31/2020 **Report Received:** 1/5/2021

(The content of this report is not confidential and may be used in communications with your organization.)

Original Project Description:

Hypothyroidism is the most common endocrine disease in dogs. A diagnosis of hypothyroidism relies on finding both appropriate clinical signs and low thyroid hormone levels. Unfortunately, other illnesses can suppress thyroid hormone levels and result in a misdiagnosis. This phenomenon of low thyroid hormone levels caused by a disease not involving the thyroid gland is known as nonthyroidal illness or euthyroid sick syndrome. It is important to distinguish between nonthyroidal illness and hypothyroidism as the treatment for each is different. Historically, the recommendation for a dog with nonthyroidal illness has been to resolve the underlying disease, followed by a recheck of thyroid hormone levels thereafter. However, the duration of time after resolution of the nonthyroidal illness necessary to perform accurate thyroid hormone level testing is unknown. This study will provide information about thyroid hormone levels during the course of nonthyroidal illness, and also establish the approximate duration of time for recovery of thyroid hormone levels to normal following illness resolution. These results will correlate clinically with more concrete recommendations for thyroid hormone level testing following resolution of nonthyroidal illness.

Publications: None at this time.

Presentations: None at this time.



Report to Grant Sponsor from Investigator:

To date, 5 cases (20% of the goal) have been enrolled.

Based on the case enrollment thus far, all dogs had normalization of total T4 by the 2-week recheck. Thus, in dogs where acute nonthyroidal illness is diagnosed, rechecking the total T4 two weeks after nonthyroidal illness resolution will result in an accurate assessment of thyroid hormone status.

As expected, 100% of dogs at the time of nonthyroidal illness diagnosis had a low total T4 and 80% had a low total T3. These values returned to normal as the disease process resolved. Free T4 was normal in all cases, as was the TSH. The TSH, in 80% of the cases, appears to significantly increase at the 2- and/or 4-week rechecks, albeit still within the reference range (future statistical analysis will determine if this is truly significant). Interestingly, in one case, the TSH increases above the reference range prior to returning to normal, a phenomenon described in humans recovering from nonthyroidal illness.

Enunciating differences between the pattern of thyroid hormone recovery in dogs and humans with naturally occurring disease will be postponed until more cases are enrolled and thus more data can be looked at to see if some of the early differences seen persist.